

Critical problems for bird conservation in the Galápagos Islands

David A. Wiedenfeld and Gustavo A. Jiménez-Uzcátegui

Received 31 May 2007; final revision accepted 14 January 2008

Cotinga 29 (2008): 22–27

La avifauna de Galápagos tiene un 50% de endemismo a nivel de especies y subespecies, típico de archipiélagos aislados. Muchas aves de Galápagos están amenazadas de extinción. Aunque varias especies se reconocen como amenazadas por UICN¹⁴, otras no son reconocidas. En Galápagos, la lista de aves amenazadas incluye 16 especies, subespecies y/o poblaciones. Entre ellas, el Pinzón de Manglar *Camarhynchus heliobates* y Cucuve de Floreana *Mimus trifasciatus* tienen poblaciones inferiores a 100 individuos. Los problemas de las especies amenazadas son la pérdida y fragmentación del hábitat, ingreso de especies introducidas que compiten por hábitat o alimento, depredadores o vectores de enfermedades, desastres naturales, pérdida de diversidad genética, actividades humanas como la pesca, agricultura y turismo sin medidas precautelares y el calentamiento global. Estas amenazas afectan a varias especies en forma específica o grupal.

The avifauna of the Galápagos Islands is quite small compared to that of an equal-sized area on the Ecuadorian mainland, with only 88 breeding or regular migrant species. However, a high proportion (c.50%) of these is endemic, either at the species or subspecies level³⁶. As is typical of island systems with a restricted land area and high endemism, many of these species are threatened with extinction.

Threats for terrestrial species are often different from those affecting marine species. For terrestrial species, threats often derive from their very limited distributions, simply as a result of the small land area of the islands, only c.45,600 km² in aggregate²⁶. Most terrestrial species do not even occur across this whole land area, but are restricted to single or a few islands, thereby having very small actual ranges. Marine species often face threats unrelated to range size, but rather to exploitation (at sea) or predation (at nesting sites).

The IUCN Red List¹⁴ includes 13 Galápagos species. The IUCN, however, does not evaluate subspecies or populations, and its taxonomy is not always the most recent. Following the up-to-date taxonomy of the South American Classification Committee of the American Ornithologists' Union²⁷ and including also Galápagos endemic subspecies results in a list of 16 taxa or populations considered threatened within the Galápagos archipelago, as will be discussed below.

Threats to Galápagos avifauna

Habitat loss. Because 96.7% of the land area of Galápagos is protected by the Galápagos National Park²⁹, habitat loss is not a significant issue for most lowland terrestrial species. However, much of the non-protected area include the most restricted, vulnerable areas in the humid highlands. These highland habitats have been disproportionately affected by agriculture, including domestic grazing,

and introduced pest plants and animals. Plant communities have been severely damaged by introduced feral goats *Capra hircus* and donkeys *Equus asinus*.

Introduced species. Introduced vertebrates are mainly predators that affect bird populations in three ways: killing adult birds (e.g., cats *Felis catus* on many species, or dogs *Canis familiaris* and pigs *Sus scrofa* on flightless or nesting species); destroying nests and young (cats, dogs, black rat *Rattus rattus* and Smooth-billed Ani *Crotophaga ani*); or competing for food (probably rats¹⁹). On the other hand, introduced plant pests affect Galápagos birds through habitat alteration, including crowding out bird food species or loss of nesting habitat. Introduced invertebrates may affect birds in two ways: competition for food or habitat alteration (for example, the cottony cushion scale *Icerya purchasi*⁶) or as parasites and vectors of diseases. These include current known threats of *Philornis downsi*, a bot-fly parasite affecting nestlings^{11,37} and Avipox³⁰. Other potential disease threats include West Nile Virus²³, avian influenza, avian malaria and others¹⁵. These diseases and parasites may kill adults or nestlings outright, make birds more susceptible to predators, or may reduce breeding success.

Natural disasters. Because of small population sizes of many of these threatened groups, natural disasters such as severe El Niño events and large volcanic eruptions may be significant to Galápagos species.

Loss of genetic diversity. Small and/or isolated populations may be threatened by inbreeding and loss of genetic diversity.

Other human activities. Fishing techniques, especially long-lining, are hazardous for many

seabirds which inadvertently get caught on hooks. Human-caused catastrophes such as petroleum spills²⁵ and other pollution or brush fires may also pose significant threats. In addition, development of land for housing, tourist facilities or agriculture can cause habitat loss. This is especially significant in the highlands of the larger islands, where native habitats are already under pressure from introduced species.

Global climate change. Possibly the most significant threat from global climate change is its potential to affect the frequency and severity of El Niño-Southern Oscillation events. El Niño may severely affect marine species, whereas the cool phase of the oscillation (La Niña events) affects terrestrial species^{32,33}. In addition, sea level rise attributable to global warming may damage coastal habitats such as mangroves and lagoons.

Species accounts

Criteria for including species, subspecies or populations in the following accounts are: 1) fewer than 1,500 individuals in the population; 2) breeding population restricted to a single island; or 3) small but unknown population and probably declining. There are 16 species, subspecies or Galápagos populations meeting at least one of these criteria. These species and the threats they face are discussed below. A summary of the threats is presented in Table 1.

Galápagos Penguin *Spheniscus mendiculus* and Flightless Cormorant *Phalacrocorax harrisi*

Because of their small population sizes, these two species (Fig. 1 for Flightless Cormorant) are threatened by natural disasters, especially El Niño events³³, loss of genetic diversity and human activity (fishing). Introduced predators, especially cats, dogs and rats may be a threat, especially to nesting penguins³².

Waved Albatross *Phoebastria irrorata*

Almost the entire population breeds on a single island, Española (Fig. 2)³. The principal threat, however, is mortality due to long-line fishing and directed fishing off the Peruvian coast²⁰. Some long-lining is permitted in the Galápagos Marine Reserve²². The species is also affected by introduced mosquitoes during the breeding season².

Galápagos Petrel *Pterodroma phaeopygia*

Threats exist both on land and at sea, where they also face long-lining as the primary threat. On land, habitat alteration by introduced plants (blackberry *Rubus* spp. and quinine trees *Cinchona pubescens*), habitat loss to development, predators, and construction of towers and powerlines threaten nesting petrels^{7,8,31}.

Greater Flamingo *Phoenicopterus ruber*

Although not globally threatened¹⁴, it is considered as threatened in Ecuador^{13,21}. The Galápagos population is quite small, between 320 and 550 individuals¹⁷. Threats include predation by pigs and other introduced species.

Galápagos Hawk *Buteo galapagoensis*

In recent years Galápagos Hawk⁴ has benefited from an increased food supply (carrión) resulting from goat eradication efforts⁵. However, it faces continued persecution by humans on inhabited islands, especially in the agricultural zones.

Galápagos Rail *Laterallus spilonotus*

The main threat to this marginally flightless species²⁴ is probably habitat loss through conversion by the introduced plants *Cinchona pubescens* (Santa Cruz) and *Rubus niveus*, but may include loss of habitat to agriculture activity as well¹², and threats from introduced predators, especially cats.

American Oystercatcher *Haematopus palliatus galapagensis*

This endemic subspecies—therefore not evaluated by IUCN¹⁴—has a very small population, probably fewer than 500 individuals, which almost certainly has never been much larger. Introduced predators may affect this ground-nesting species.

Lava Gull *Leucophaeus fuliginosus*

The current population of Lava Gull (Fig. 3) is probably smaller than the 800 individuals estimated by Snow & Snow²⁸ in the late 1960s. An estimate for southern Santa Cruz Island¹, where one of the larger populations exists, resulted in 81 individuals. The IUCN¹⁴ lists the species as Vulnerable, which probably understates the seriousness of its status.

Vermilion Flycatcher *Pyrocephalus rubinus nanus* and *P. r. dubius*

Populations of the two Galápagos subspecies, *P. r. nanus* and *P. r. dubius*, are quite small and apparently declining; the species is already extinct on San Cristóbal³⁶. However, the reasons for the decline are unclear.

Galápagos Martin *Progne modesta*

This is possibly the least-known species in Galápagos. Its population is very small, probably much fewer than 500 individuals and the species can only be found consistently at a few sites³⁶. It is listed by the IUCN¹⁴ as Vulnerable, but the species' scarcity and probable decline might warrant Endangered status. Threats may include introduced diseases or parasites, and possibly introduced predators such as rats.

Table I. List of threats. Known threats are indicated by a *; those threats not well known but suspected are marked '?'. Threat category from IUCN¹⁴ CR (Critically Endangered), EN (Endangered), VU (Vulnerable), LC (Least Concern).

Species	IUCN Status	Introduced vertebrates	Introduced plants	Introduced invertebrates	Diseases or parasites	Habitat loss	Natural disasters	Global climate change	Loss of genetic diversity	Other human activity
<i>Spheniscus mendiculus</i>	EN	Cats, dogs, rats on nesting birds ³³			?	Volcanic eruptions, especially on Fernandina	Especially El Niño events ³³	•	•	Fishing, through tangling in nets ³⁴ , and competition
<i>Phoebastria irrorata</i>	CR				•		•	•		Long-line and directed fishing ²⁰
<i>Pterodroma phaeopygia</i>	CR	Cats, pigs, and rats on eggs and nestlings	<i>Cinchona pubescens</i> and <i>Rubus</i> spp. over-grow colonies ³¹			•				Collisions with towers and powerlines ^{7,8} , potential wind power development on Santa Cruz
<i>Phalacrocorax harrisi</i>	EN	Predators on nests and nestlings ³²				Volcanic eruptions on Fernandina	Especially El Niño events ³²	•	•	Fishing, through tangling in nets, and competition
<i>Phoenicopterus ruber</i>	LC	Predators on nests, especially pigs			Die-offs possibly caused by avian cholera	•	•	Lagoons may be threatened by sea level rise		•
<i>Buteo galapagoensis</i>	VU				•				Small populations on some islands ⁴	Persecution on inhabited islands in agricultural zones
<i>Laterallus spilonotus</i>	VU	Especially cats ³⁴ , but also rats	<i>Cinchona pubescens</i> on Santa Cruz; <i>Rubus niveus</i>			Continuing conversion of highland habitat to agriculture ¹²			?	
<i>Haematopus palliatus galapagensis</i>	none	Predators on nests							?	
<i>Leucophaeus fuliginosus</i>	VU	Predators on nests, especially cats and dogs				?			?	Fishing activities, hooks and nets. Feed on refuse in town.
<i>Pyrocephalus rubinus nanus</i> and <i>P. r. dubius</i>	none	?		?	Possibly avian pox or <i>Philornis downsi</i> ³⁷	?			?	
<i>Progne modesta</i>	VU	Rats?			Avian pox or <i>Philornis downsi</i> ?	?		?	?	
<i>Mimus trifasciatus</i>	EN	Predators, chiefly rats, but not yet present on islets where the species occurs; <i>Crotophaga ani</i> ³⁵		?	<i>Philornis downsi</i> ³⁷ , pox ³	?	Prolonged drought from La Niña events	•	Divided into two very small populations on separate islets	
<i>Mimus melanotis</i>	EN	Probably cats and rats	?	?	<i>Philornis downsi</i> ³⁷ and avian pox	?	•	?		?
<i>Mimus macdonaldi</i>	VU			?	?		Prolonged drought from La Niña events			•
<i>Camarhynchus pauper</i>	VU	Predators such as cats and rats	?	?	<i>Philornis downsi</i> ³⁷	?			?	
<i>Camarhynchus heliobates</i>	CR	Predators such as cats, rats, <i>Crotophaga ani</i>		Fire ants kill nestlings	<i>Philornis downsi</i> and avian pox	•	Prolonged drought from La Niña events	Sea level rise threatens mangroves	Very small, isolated populations	•



Figure 1. Flightless Cormorant *Phalacrocorax harrisi* on nest, Cape Douglas, Fernandina Island, Galápagos, 8 September 2002 (David A. Wiedenfeld)



Figure 2. Waved Albatross *Phoebastria irrorata*, Cevallos Point, Española Island, Galápagos, 1 June 2003 (David A. Wiedenfeld)



Figure 3. Lava Gull *Leucophaeus fuliginosus* on nest, west of Puerto Ayora, Santa Cruz Island, Galápagos, 6 May 2005 (David A. Wiedenfeld)



Figure 4. Floreana Mockingbird *Mimus trifasciatus*, Champion Island, Galápagos, 21 May 2004 (David A. Wiedenfeld)

Floreana Mockingbird *Mimus trifasciatus*

The Floreana Mockingbird (Fig. 4) probably numbers no more than 150 individuals, and the most recent surveys revealed a sharp decline in 2006¹⁶ followed by a partial recovery in 2007¹⁸. It is now confined to two small islets. The accidental introduction of predators must be avoided as they could deplete its small population rapidly. The introduced fly parasite *Philornis downsi* is known from both islets³⁷, whilst avian pox has also been recorded⁹. Other potential threats include introduced Smooth-billed Ani *Crotophaga ani*, prolonged droughts and inbreeding as there is apparently no gene flow between the two islets.

San Cristóbal Mockingbird *Mimus melanotis*

The population of the San Cristóbal Mockingbird is unknown, but is at most only a few thousand birds, all restricted to one island.

Española Mockingbird *Mimus macdonaldi*

The species is restricted to a single island, but one with no introduced mammalian predators¹⁹, and *Philornis downsi* has apparently not yet reached the island³⁷. Because Española has no consistent fresh water, prolonged drought can cause significant mortality.

Medium Tree-Finch *Camarhynchus pauper*

The population size of the Medium Tree-Finch is unknown but probably very small, as the species is restricted to Floreana Island. IUCN lists the species as Vulnerable, but its small population size and significant threats suggest upgrading it to Endangered.

Mangrove Finch *Camarhynchus heliobates*

The Mangrove Finch is one of the most endangered species in the world. Its population is fewer than 100 individuals, possibly as few as 50, restricted to only two sites with a total of c.12 ha on Isabela Island¹⁰. With a population size so small, the

Mangrove Finch faces threats from nearly everything; avian diseases and introduced predators have been reported, whilst catastrophic events may prove disastrous.

Discussion

Although the Galápagos Islands have to date lost no bird species to extinction³⁸, a significant percentage of the Galápagos avifauna is threatened²¹. The main threats are introduced mammalian predators, parasites, diseases and invasive plants. The combination of very small populations of many Galápagos species (several probably never occurred in very large numbers) combined with their oceanic-island naiveté³⁸ means Galápagos birds often have a perilous population status. Even though the Galápagos National Park Service and Charles Darwin Foundation have made strong conservation efforts in the Galápagos, problems remain and in many cases the threats have grown¹². Species with such restricted ranges and small populations can never be assumed to be beyond threat. Conservationists must continue their efforts to solve the difficult and thorny problems to protect such unique species as those found in the Galápagos Islands.

Acknowledgements

We thank all of the wardens of the Galápagos National Park Service, volunteers, scholarship students and personnel of the Charles Darwin Research Station, and scientists of various universities, who between them have helped acquire information on the birds of Galápagos and the threats they face. Galápagos Conservancy, Friends of Galápagos–Switzerland, Galápagos Conservation Foundation, GEF Project ECU/00/G31, St Louis Zoo, University of Missouri, Sea World, the Japan Embassy, Swarovski and others have financed studies of birds; without their support it would not have been possible to complete this study. We also express our appreciation to the Galápagos National Park Service for allowing us to work on the islands, and to the Charles Darwin Foundation and Charles Darwin Research Station for logistical support. Juan Freile and Tatiana Santander helped to improve the manuscript.

References

1. Aguirre, D. (2007) Aspectos etológicos, influencia antropogénica y distribución de la Gaviota de Lava (*Larus fuliginosus*), una especie endémica del archipiélago de Galápagos. Tesis de Licenciatura. Quito: Univ. Central de Ecuador.
2. Anderson, D. J. & Fortner, S. (1988) Waved Albatross egg neglect and associated mosquito ectoparasitism. *Condor* 90: 727–729.
3. Anderson, D. J., Huyvaert, K. P., Apanius, V., Townsend, H., Gillikin, C. L., Hill, L. D., Juola, F., Porter, E. T., Wood, D. R., Loughheed, C. & Vargas, H. (2002) Population size and trends of the Waved Albatross *Phoebastria irrorata*. *Marine Orn.* 30: 63–69.
4. Bollmer, J. L., Kimball, R. T., Whiteman, N. K., Sarasola, J. H. & Parker, P. G. (2006) Phylogeography of the Galápagos Hawk (*Buteo galapagoensis*): a recent arrival to the Galápagos Islands. *Mol. Phyl. & Evol.* 39: 237–247.
5. Campbell, K., Donlan, J., Cruz, F. & Carrión, V. (2002) Eradication of feral goats *Capra hircus* from Santa Cruz Island, Galápagos, Ecuador. *Oryx* 38: 328–333.
6. Causton, C. E., Peck, S. B., Sinclair, B. J., Roque-Albelo, L., Hodgson, C. J. & Landry, B. (2006) Alien insects: threats and implications for conservation of Galápagos Islands. *Ann. Ent. Soc. Amer.* 99: 121–143.
7. Cruz-Delgado, F. (2006) Estudio ornitológico del Petrel de Galápagos para el proyecto eólico en Santa Cruz y Baltra. In: Jiménez-Uzcátegui, G. (ed.) Estudio biológico en Santa Rosa, El Camote, Isla Santa Cruz previo a la colocación del parque eólico. Unpubl. rep. Puerto Ayora: Fundación Charles Darwin, ERGAL & Servicio Parque Nacional Galápagos.
8. Cruz-Delgado, F. & Wiedenfeld, D. A. (2005) Report of results of the ornithological study of the Galápagos Petrel on Cerro Tropezón, San Cristóbal Island, July–October 2004 and February–March 2005. Unpubl. rep. Puerto Ayora: Fundación Charles Darwin, e7 Fund & Servicio Parque Nacional Galápagos.
9. Curry, R. L. (1985) Breeding and survival of Galápagos mockingbirds during El Niño. In: Robinson, G. & del Pino, E. M. (eds.) *El Niño en las Islas Galápagos: el evento de 1982–1983*. Quito: Fundación Charles Darwin.
10. Dvorak, M., Vargas, H., Fessl, B. & Tebbich, S. (2004) On the verge of extinction: a survey of the Mangrove Finch *Cactospiza heliobates* and its habitat on the Galapagos Islands. *Oryx* 38: 1–9.
11. Fessl, B. & Tebbich, S. (2002) *Philornis downsi*—a recently discovered parasite on the Galápagos archipelago—a threat for Darwin's finches? *Ibis* 144: 445–451.
12. Gibbs, J. P., Shriver, W. G. & Vargas, H. (2003) An assessment of a Galapagos Rail population over thirteen years (1986–2000). *J. Field Orn.* 74: 136–140.
13. Granizo, T., Pacheco, C., Ribadeneira, M. B., Guerrero, M. & Suárez, L. (2002) *Libro rojo de las aves del Ecuador*. Quito: SIMBIOE, Conservation International, EcoCiencia, Ministerio del Ambiente & IUCN.
14. IUCN (2007) 2007 IUCN Red List of Threatened Species. Available at: www.iucnredlist.org.
15. Jiménez-Uzcátegui, G. (2002) Enfermedades bacterianas, víricas, micóticas y parasitarias que posiblemente pueden afectar a las aves silvestres de Galápagos. Unpubl. rep. Puerto Ayora: Fundación Charles Darwin & Servicio Parque Nacional Galápagos.
16. Jiménez-Uzcátegui, G. (2006) Censo de Cucuve de Floreana *Nesomimus trifasciatus* 2006. Unpubl.

- rep. Puerto Ayora: Fundación Charles Darwin & Servicio Parque Nacional Galápagos.
17. Jiménez-Uzcátegui, G. (2006) Censo parcial de Flamencos *Phoenicopterus ruber* 2006. Unpubl. rep. Puerto Ayora: Fundación Charles Darwin & Servicio Parque Nacional Galápagos.
 18. Jiménez-Uzcátegui, G. (2007) Censo de Cucuve de Floreana *Nesomimus trifasciatus* 2007. Unpubl. rep. Puerto Ayora: Fundación Charles Darwin & Servicio Parque Nacional Galápagos.
 19. Jiménez-Uzcátegui, G., Carrión, V., Zabala, J., Buitrón, P. & Milstead, B. (2007) Estado de vertebrados introducidos en Galápagos. In: *Informe Galápagos 2006-2007*. Puerto Ayora: Fundación Charles Darwin, Servicio Parque Nacional Galápagos & INGALA.
 20. Jiménez-Uzcátegui, G., Mangel, J., Alfaro-Shigueto, J. & Anderson, D. J. (2006) Fishery bycatch of the Waved Albatross *Phoebastria irrorata*, a need for implementation of agreements. *Galápagos Res.* 64: 7-9.
 21. Jiménez-Uzcátegui, G., Milstead, B., Márquez, C., Zabala, J., Buitrón, P., Llerena, A., Salazar, S. & Fessl, B. (2007) Vertebrados de Galápagos: estado de amenaza y acciones de conservación. In: *Informe Galápagos 2006-2007*. Puerto Ayora: Fundación Charles Darwin, Servicio Parque Nacional Galápagos & INGALA.
 22. Junta de Manejo Participativo (2005) *Boletín informativo de la Junta de Manejo Participativo de la Reserva Marina de Galápagos, 22 diciembre 2005*. Puerto Ayora: Junta de Manejo Participativo.
 23. Kilpatrick, A. M., Daszak, P., Goodman, S., Rogg, H., Kramer, L. D., Cedeño, V. & Cunningham, A. (2006) Predicting pathogen introduction: West Nile virus spread to Galápagos. *Conserv. Biol.* 20: 1224-1231.
 24. Livezey, B. C. (2003) Evolution of flightlessness in rails (Gruiformes: Rallidae): phylogenetic, ecomorphological, and ontogenetic perspectives. *Orn. Monogr.* 53. Washington DC: American Ornithologists' Union.
 25. Loughheed, C. W., Edgar, G. J. & Snell, H. L. (2001) Biological impacts of the Jessica oil spill on the Galápagos environment: preliminary report. Unpubl. rep. Puerto Ayora: Fundación Charles Darwin.
 26. Perry, R. (1984) The islands and their history. In: Perry, R. (ed.) *Key environments Galapagos*. Oxford: Pergamon Press.
 27. Remsen, J. V., Cadena, C. D., Jaramillo, A., Nores, M., Pacheco, J. F., Robbins, M. B., Schulenberg, T. S., Stiles, F. G., Stotz, D. F. & Zimmer, K. J. (2007) A classification of the bird species of South America. Available at: www.museum.lsu.edu/~Remsen/SACCBaseline.html.
 28. Snow, B. K. & Snow, D. W. (1967) Observations of the Lava Gull *Larus fuliginosus*. *Ibis* 111: 30-35.
 29. SPNG (2007) Ubicación geográfica. Parque Nacional Galápagos Ecuador. Available at: www.galapagospark.org/png/interna.php.
 30. Thiel, T., Whiteman, N., Tirapé, A., Baquero, M. I., Cedeño, V., Walsh, T., Jiménez-Uzcátegui, G. & Parker, P. (2005) Identification of canary pox-like viruses infecting endemic birds in the Galapagos Islands. *J. Wildl. Dis.* 41: 342-353.
 31. Valarezo, J. C. (2006) Redescubriendo el hábitat del Petrel de Galápagos (*Pterodroma phaeopygia*). *Bol. Soc. Antioqueña Orn.* 16: 70-79.
 32. Vargas, H. (2006) The ecology of small populations of birds in changing climate. PhD dissertation. Oxford: University of Oxford.
 33. Vargas, H., Harrison, S., Rea, S. & Macdonald, D. W. (2006) Biological effects of El Niño on the Galapagos Penguin. *Biol. Conserv.* 127: 107-114.
 34. Vinuesa, C. L. & Wiedenfeld, D. A. (2003) Censo de pingüinos en Puerto Villamil-Isabela. Unpubl. rep. Puerto Ayora: Fundación Charles Darwin.
 35. Wiedenfeld, D. A. (2003) Floreana Mockingbird survey 2003. Unpubl. rep. Puerto Ayora: Fundación Charles Darwin & Servicio Parque Nacional Galápagos.
 36. Wiedenfeld, D. A. (2006) The avifauna of the Galapagos Islands, Ecuador. *Checklist* 2: 1-27.
 37. Wiedenfeld, D. A., Jiménez-Uzcátegui, G. A., Fessl, B., Kleindorfer, S. & Valarezo, J. C. (2007) Distribution of the introduced parasitic fly *Philornis downsi* (Diptera, Muscidae) in the Galapagos Islands. *Pacific Conserv. Biol.* 13: 14-19.
 38. Wikelski, M., Foufopoulos, J., Vargas, H. & Snell, H. (2004) Galápagos birds and diseases: invasive pathogens as threats for island species. *Ecol. Soc.* 9: 5.

David A. Wiedenfeld

American Bird Conservancy, PO Box 249, The Plains, Virginia 20198, USA. E-mail: dwiedenfeld@abcbirds.org.

Gustavo A. Jiménez-Uzcátegui

Estación Científica Charles Darwin, Puerto Ayora, Galápagos, Ecuador. E-mail: gjimenez@fcdarwin.org.ec.