

Altitudinal and geographical range extension for Bicoloured Antvireo *Dysithamnus occidentalis punctitectus* in south-east Ecuador, with notes on its nesting ecology

J. Berton C. Harris, Rolando L. Carpio A., Mary K. Chambers and Harold F. Greeney

Received 23 September 2007; final revision accepted 10 March 2008

Cotinga 30 (2008): 63–65

Presentamos una extensión del rango altitudinal y geográfico para el Batarito Bicolor *Dysithamnus occidentalis punctitectus* en la Cordillera de Sabanilla, parroquia Valladolid, provincia de Zamora-Chinchipec, Ecuador. En abril y mayo de 2007 en la Reserva Biológica Tapichalaca en un rango de 2.300–2.460 msnm, se observó en tres ocasiones y capturó en redes de neblina el 11 de junio a *D. occidentalis*. Estos registros extienden el rango altitudinal de la especie en Ecuador 260 m y el rango geográfico 160 km al sur-oeste en Ecuador. Adicionalmente, presentamos nueva información sobre la ecología de anidación y comportamiento reproductivo de la especie del oriente al Ecuador (provincia de Napo).

Bicoloured Antvireo *Dysithamnus occidentalis* is an inconspicuous, low-density bird that occurs locally on the west slope of the Andes in western Colombia^{2–6,15} and northern Ecuador, and disjunctly along the east slope of the Ecuadorian Andes and in northern Peru^{1,2,9,11–15}. It was essentially unknown in life until 1991¹⁴ and is considered Vulnerable due to habitat loss^{2,4,11,15}. Two subspecies are recognised. *D. o. occidentalis* occurs in Colombia at 900–2,800 m^{4–6} and in north-west Ecuador at 2,200 m⁹ (subspecies inferred by range). *D. o. punctitectus* is known from Ecuador's east slope at 1,500–2,050 m¹¹, including specimens taken in the 1920s 'below Oyacachi', 'reportedly near Baeza', and 'Sumaco abajo'¹¹, and two recent specimens, from 1,500 m, at Río Abanico near Volcán Sangay⁹. Even more recently, the species was recorded further south, on the west slope of the Cordillera del Cóndor near San Pedro de Apondios, prov. Morona-Santiago, at 1,600–1,900 m¹. It is also now known at 2,000–2,500 m in Peru, on the south slope of the Cordillera del Cóndor, dpto. Cajamarca, and near Abra Patricia, south of the río Marañón, dpto. San Martín¹³.

The species' reproductive biology is still poorly known. Only two nests have been described^{7,8} and data on eggs and incubation behaviour are available from just one nest⁹.

Here we report *D. o. punctitectus* (inferred by range) from the Cordillera de Sabanilla, prov. Zamora-Chinchipec, Ecuador, thereby extending the species' known altitudinal range in Ecuador by 260 m (and that of *punctitectus* by 410 m) and the geographical range by 160 km south-west from the Cordillera del Cóndor¹. We also present further observations on nesting ecology and behaviour from prov. Napo, north-east Ecuador (at Yanayacu Biological Station; 1,950–2,100 m; 00°36'S 77°53'W).

Range extension

We observed Bicoloured Antvireos in mature forest at 2,300–2,460 m (04°29'S 79°07'W), in Tapichalaca Biological Reserve, a 2,870-ha protected area administered by Fundación Jocotoco, above the town of Valladolid. Forest in this area, described as upper subtropical forest¹⁰, has a mean canopy height of c.10 m, with 20-m emergent crowns, and receives c.4 m of rainfall p.a. The canopy is characterised by Moraceae (*Ficus* sp.), Euphorbiaceae (*Croton* sp.), Lauraceae and Rubiaceae, and the understorey is largely comprised of *Chusquea* sp. bamboo (Poaceae), Piperaceae and Melastomataceae. Steep slopes and heavy epiphyte loads make the forest prone to landslides and treefalls. As reported earlier^{1,5,14}, we encountered antvireos in areas of localised early-successional habitat such as bamboo thickets and vine tangles, in otherwise undisturbed forest.

Whilst mist-netting between 25 April and 14 June 2007 we observed Bicoloured Antvireos four times and captured two individuals. On 25 April, RLCA observed a closely associated group of one male and two females, for ten minutes. On 1 May RLCA and JBCH observed and made sound-recordings (to be archived at the Macaulay Library, Cornell University) of two males and two females, for 15 minutes. On 7 May RLCA observed two males for five minutes and on 8 May RLCA observed a pair for two minutes. On 11 June we mist-netted a presumed pair in nets 25 m apart. We caught the female 30 minutes after capturing the male. We estimated skull ossification to be 100% for both individuals and the female had a receding brood patch.

All individual antvireos remained within 2 m of each other during the observations and we never observed an agonistic interaction. They made frequent but quiet vocalisations of both the smooth *peeu* and fast scold *jeer-deer-dur* types¹¹, corroborat-

ing Greeney's⁷ observation that the species is very vocal yet inconspicuous because the calls are so muted. We never observed antvireos forage within a mixed-species flock at Tapichalaca. They never foraged more than 2 m above ground and usually at less than 1 m, or on the ground, as described by Whitney¹⁴, contrary to the brief observations of Ágreda *et al.*¹. All of our observations involved at least two individuals and twice we observed 3+ individuals. Our record of a male and two females might have represented a family group, although all appeared to be adults¹³. It is unclear, however, why the two apparent pairs we observed remained within 2 m of each other for 15 minutes without exhibiting agonistic behaviour.

D. occidentalis is an apparently rare resident at Tapichalaca Biological Reserve. Despite that Tapichalaca has been frequented by experienced observers since 1998, *D. occidentalis* was not definitely recorded until 2007. That we always observed at least two individuals, and the presence of a brood patch on the captured female, suggest that the species is resident in the region. The species' apparent scarcity is partially explained by its inconspicuous behaviour and quiet vocalisations, but even when these factors are considered, we suggest a total population size of <30 individuals in the reserve. Recent range extensions in Ecuador^{19,13} suggest that *D. occidentalis* may yet be discovered in mature forest at 1,500–2,500 m in Podocarpus National Park, to the north of Tapichalaca, or in the Cordillera de las Lagunillas to the south.

Nesting ecology and behaviour

At Yanayacu *D. occidentalis*, whilst frequently found as solitary pairs, often joins small understorey flocks comprised of Spotted Barbtails *Premnoplex brunnescens*, Grey-breasted Wood Wrens *Henicorhina leucophrys* and Chestnut-capped Brush Finches *Buarremon brunneinucha*. Nesting has previously been documented in November and December from the area^{7,8}. Here we present data from four other active and three unoccupied nests. We found nests under construction in early March 2003 and late October 2006. We also found a nest with incubation underway in mid-November 2002 and one with two older nestlings in early December 2004. In the same area, R. A. Gelis observed a juvenile with two adults in mid-August 2003. Clutch size at all nests was two eggs. Eggs at one nest measured 21.5 × 16.3 and 21.5 × 16.4 mm, and those at a second 22.1 × 16.2 and 22.2 × 16.4 mm. Using previous egg measurements from this area⁸ we calculate mean (± SD) dimensions as 21.9 ± 0.3 × 16.4 ± 0.1 mm. Nests in the area, including previously published data^{7,8}, were found in small (mean dbh 14 ± 5 mm) saplings of the following families (numbers of nests in parentheses):

Solanaceae (4), Piperaceae (1), Melastomataceae (1), Myristicaceae (1), and unknown (2). Mean substrate height was 2.3 ± 0.3 m and mean nest height 1.5 ± 0.4 m. Nests were 0–25 m from small streams (mean 6.4 ± 10.1), always in mature forest, but usually in areas of natural disturbance such as treefalls. All nests were situated next to the trunk of the supporting tree and suspended between two thin (mean 4.6 ± 1.6 mm diameter) horizontal branches, on average separated vertically by 3.9 ± 1.3 cm. Mean nest dimensions (cm) were: outer diameter 9.6 ± 1.0; outer height 7.2 ± 0.8; inner diameter 6.6 ± 0.7; and inner depth 5.2 ± 0.9. All nests were dense pendant cups woven entirely of dark rootlets.

In sum, records from Yanayacu suggest breeding occurs year-round in this area, with a fairly defined peak towards the late drier season, in November–December.

Acknowledgements

Fundación Jocotoco and the World Land Trust provided funding and logistical support. We thank the Fundación Jocotoco office staff and Tapichalaca park guards, as well as our dedicated field assistants, L. Reid, L. Marshall and M. Wickens. A. Ágreda, D. Haskell, N. Krabbe, J. Phillips, R. S. Ridgely and B. Scheffers made helpful comments on the manuscript. We also thank P. Álvarez, G. Budney, N. Hollingshead, M. Juiña, D. F. Lane, T. S. Schulenberg and F. Sornoza. HFG acknowledges the ongoing support of John V. & the late Ruth Ann Moore. This is publication no. 131 of the Yanayacu Natural History Research Group.

References

1. Ágreda, A., Nilsson, J., Tonato, L. & Román, H. (2005) Range extension for, and description of the juvenile of, Bicoloured Antvireo *Dysithamnus occidentalis punctitectus* in Ecuador. *Cotinga* 24: 20–21.
2. BirdLife International (2000) *Threatened birds of the world*. Cambridge, UK: BirdLife International & Barcelona: Lynx Edicions.
3. Bond, J. & Meyer de Schauensee, R. (1940) On some birds from southern Colombia. *Proc. Acad. Nat. Sci. Philadelphia* 92: 153–169.
4. Collar, N. J., Gonzaga, L. P., Krabbe, N., Madroño Nieto, A., Naranjo, L. G., Parker, T. A. & Wege, D. C. (1992) *Threatened birds of the Americas: the ICBP/IUCN Red Data book*. Cambridge, UK: International Council for Bird Preservation.
5. Donegan, T. M. & Dávalos, L. M. (1999) Ornithological observations from Reserva Natural Tambito, Cauca, south-west Colombia. *Cotinga* 12: 48–55.
6. Echeverry-Galvis, M. Á. & Córdoba-Córdoba, S. (2007) New distributional and other bird records from Tatamá Massif, West Andes, Colombia. *Bull. Brit. Orn. Club* 127: 213–224.
7. Greeney, H. F. (2002) First description of the nest for Bicoloured Antvireo (*Dysithamnus occiden-*

- talís*), with notes on its behavior in eastern Ecuador. *Orn. Neotrop.* 13: 1–3.
8. Greeney, H. F. (2004) Breeding behavior of the Bicolored Antvireo (*Dysithamnus occidentalis*). *Orn. Neotrop.* 15: 349–356.
 9. Krabbe, N. & Palacio, J. (1999) Range extensions of Bicoloured Antvireo *Dysithamnus occidentalis* in Ecuador. *Cotinga* 11: 48.
 10. Krabbe, N., Agro, D. J., Rice, N. H., Jácome, N., Navarrete, L. & Sornoza M., F. (1999) A new species of antpitta (Formicariidae: *Grallaria*) from the southern Ecuadorian Andes. *Auk* 116: 882–890.
 11. Ridgely, R. S. & Greenfield, P. J. (2001) *The birds of Ecuador*, 1. Ithaca, NY: Cornell University Press.
 12. Ridgely, R. S. & Tudor, G. (1994) *The birds of South America*, 2. Austin: University of Texas Press.
 13. Schulenberg, T. S., Stotz, D. F., Lane, D. F., O'Neill, J. P. & Parker, T. A. (2007) *Birds of Peru*. Princeton, NJ: Princeton University Press.
 14. Whitney, B. M. (1992) Observations on the systematics, behavior, and vocalizations of “*Thamnomanes*” *occidentalis* (Formicariidae). *Auk* 109: 302–308.
 15. Zimmer, K. J. & Isler, M. L. (2003) Family Thamnophilidae (typical antbirds). In: del Hoyo, J., Elliott, A. & Christie, D. A. (eds.) *Handbook of the birds of the world*, 8. Barcelona: Lynx Edicions.

J. Berton C. Harris, Rolando L. Carpio A. and Mary K. Chambers

Fundación de Conservación Jocotoco, Av. Los Shyris N37-146 y El Comercio, Quito, Ecuador. E-mail: bertdichrozona@hotmail.com.

Harold F. Greeney

Yanayacu Biological Station & Center for Creative Studies, Cosanga, Ecuador, c/o 721 Foch y Amazonas, Quito, Ecuador.