

### First documented record of Veery *Catharus fuscescens* in southern Brazil

Parts of Amazonia, the eastern Andes and sometimes even Central America have been considered within the winter range of Veery *Catharus fuscescens*. However, Remsen<sup>3</sup> demonstrated that most winter records are from north and centre-west Brazil. Further south, it has been recorded in São Paulo, by Pereira *et al.*<sup>2</sup>, who collected a male in 1985 and a female in 1986, in Salesópolis and Iguape municipalities, respectively, whilst Willis & Oniki<sup>7</sup> and E. O. Willis *in Remsen*<sup>3</sup> observed 13 individuals, on 12 dates, at altitudes of 250–1,000 m, between late November–early March. In Rio de Janeiro, there is a record by D. F. Stotz in January 1992, at Praia Seca, and another by T. Sigris, in November, at Tinguá<sup>6</sup>. The southernmost record is from Serranía San Luis National Park, in Paraguay<sup>5</sup>.

On 11 February 2009, during an ornithological inventory of Fazenda Santa Alice, Rio Negrinho municipality, Santa Catarina (26°29'22.7"S 49°29'40.3"W) a *C. fuscescens* was mist-netted at c.09h00. It was a first-year, sex unknown, and was ringed, measured and photographed (Fig. 1). Its measurements were as follows: bill (exposed culmen) 14.0 mm; tarsus 24.8 mm; wing (max.) 90.0 mm; tail 68.3 mm; total length 175.0 mm; mass 30 g. The bird was not moulting any feathers.

Fazenda Santa Alice lies at 800–1,100 m<sup>1,4</sup>. The area where the Veery was trapped is a *capoeira* of *Baccharis dracunculifolia*, with many pioneer species such as *Myrsine coriacea*, *Schinus therebentifolius*, *Zanthoxylum rhoifolium*, *Rhamnus sphaeosperma*, *Matayba*



Figure 1. Veery *Catharus fuscescens*, Fazenda Santa Alice, Santa Catarina, 11 February 2009 (Nicholas Kaminski)

*elaegnoides*, *Drymis brasiliensis* and some Melastomataceae. The general environment is a matrix of primary forest and much old secondary forest, but dominated by a vast monoculture of *Pinus taeda*.

This record brings not only an expansion of the area of wintering (*sensu* Remsen)<sup>3</sup>, but also a new vegetation typology with maximum and minimum temperatures very different from Amazonia and the Cerrado<sup>3</sup>, and even from the Atlantic Forest<sup>7</sup>. However, only with further observations, and preferably continued use of mist-nets, will it be possible to determine if *C. fuscescens* is regular in this region of Brazil.

#### Acknowledgements

Battistella Florestal funded the work. Glauco Kohler made suggestions concerning the project, Carlos Alberto Borchardt Jr. helped identify the bird, and André Ferrari translated the manuscript.

#### References

1. Klein, M. R. (1978) Mapa fitogeográfico do Estado de Santa Catarina. *Flora Ilustrada Catarinense* 5: 1–24.
2. Pereira, L. E., Sachetta, L. A. & Souza, D. M. (1988) *Catharus fuscescens* (Stephens, 1817), nova espécie para o Estado de São Paulo. *Anais III Encontro Nac. Anilhadores Aves (ENAV)*: 111–112.
3. Remsen, J. V. (2001) True winter range of the Veery (*Catharus fuscescens*): lessons for determining winter ranges of species that winter in the tropics. *Auk* 118: 838–848.
4. Roderjan, C. V., Galvão, F., Kuniyoshi, Y. S. & Hatschbach, G. G. (2000) As unidades fitogeográficas do Estado do Paraná. *Ciência e Ambiente* 24: 75–92.
5. Robbins, M. B., Faucett, R. C. & Rice, N. H. (1999) Avifauna of a Paraguayan cerrado locality: Parque Nacional Serranía San Luis, depto. Concepción. *Wilson Bull.* 111: 216–228.
6. Sigris, T. (2006) *Aves do Brasil: uma visão artística*. São Paulo: Ed. Avis Brasilis.
7. Willis, E. O. & Oniki, Y. (1993) New and reconfirmed birds from the state of São Paulo, Brazil, with notes on disappearing species. *Bull. Brit. Orn. Club* 113: 23–34.

#### Nicholas Kaminski

Alberto Erthal, 446 São Lourenço, Curitiba CEP 82.210-200, Paraná, Brazil. E-mail: nicholas.kaminski@yahoo.com.br.

Received 6 July 2009; final revision accepted 1 July 2010 (published online 16 March 2011)