

First records of Blue-billed Black Tyrant *Knipolegus cyanirostris* for Goiás, Brazil

Blue-billed Black Tyrant *Knipolegus cyanirostris* is usually found in humid and gallery forests, principally in southern and south-eastern Brazil, as well as the central west (Mato Grosso do Sul) and parts of Argentina, Paraguay and Uruguay⁵⁻⁷. In Brazil, it is considered migratory in the states



Figure 1. Male Blue-billed Black Tyrant *Knipolegus cyanirostris* mist-netted in the municipality of Catalão, south-east Goiás, Brazil, 10 August 2008 (Sandro Barata Berg)

of Espírito Santo, Rio de Janeiro⁷ and Mato Grosso do Sul⁶.

The only known record in central Brazil is a specimen collected on 1 July 1963 deposited at the Museu de Zoologia da Universidade de São Paulo (MZUSP 54764) from Planaltina, Distrito Federal (DF)⁹. Here, we report the first records in Goiás^{2,4}.

During regional surveys of the Serra do Facão Hydroelectric Dam, in the rio São Marcos basin, in south-east Goiás, we twice found individuals of *K. cyanirostris*. Vegetation is typical of the Cerrado, with a mosaic of dry forest and gallery forest, savanna (*cerrado sensu stricto*) and grasslands (*campo sujo* and *campo limpo*), as well as pastures and plantations (especially soya, bean and *Eucalyptus*), bogs, rivers and degraded areas. Description of the Cerrado's physiognomies follows Oliveira-Filho & Ratter³.

On 10 August 2008 a male Blue-billed Black Tyrant was mist-netted (Fig. 1) in a gallery forest (17°56'22"S 47°39'33"W; 750 m) flanking a small affluent of the rio São Marcos, Pires Belo district, Catalão municipality. It had red irides, a bluish-grey bill and black plumage. The bird was collected and deposited in the Coleção Ornitológica Marcelo Bagno (COMB 2961) at the Universidade de Brasília. On 16 August 2008, we observed a pair in scrub at

the edge of dry forest (17°45'02"S 47°41'45"W; 772 m).

These records extend the range of *K. cyanirostris* to central Brazil (Fig. 2) and 260 km south-west (from the Distrito Federal) and 130 km north (from the Triângulo Mineiro, Minas Gerais). Occasional records of other *Knipolegus* spp. in localities distant from their main ranges are also available^{1,7,9}, probably because of their partially migratory habits^{7,8}. Thus, the occurrence of Blue-billed Black Tyrant in south-east Goiás is expected, although further data are needed to understand its spatial and temporal occurrence there.

Acknowledgements

We are especially grateful to Nathália Machado e Sousa for her collaboration and company during our visit to Goiânia. José Hidasí (Fundação Museu Ornitológico de Goiânia) and Luís Fábio Silveira (MZUSP) permitted us to examine specimens. Aline Henrique Côrrea provided additional information concerning the specimen in MZUSP, and Miguel Marini (COMB) also offered assistance. Marcos Robalinho Lima helped us prepare an English version of the manuscript. Marcelo Vasconcelos and Guy Kirwan also made substantial contributions to the submitted version. CEMAVE and IBAMA granted permits to catch and collect (no. 3043/1) birds. We are also grateful to Serra do Facão Energia S.A. and Fundação Pró-Natureza (FUNATURA) for logistical support.

References

1. Bornschein, M. R., Cáceres, N. C., Ferreira, W. V., Freitas, D. R. & Pichorim, M. (2003) Primeiro registro de *Knipolegus aterrimus* Kaup, 1853 no Brasil (Tyrannidae). *Ararajuba* 11: 131–145.
2. Hidasí, J. (2007) *Aves de Goiás*. Goiânia: Ed. Universidade Católica de Goiás.
3. Oliveira-Filho, A. T. & Ratter, J. A. (2002) Vegetation physiognomies and woody flora of the Cerrado biome. In: Oliveira, P. S. & Marquis,




Figure 2. Range of Blue-billed Black Tyrant *Knipolegus cyanirostris* based on Ridgely & Tudor⁶. The black asterisk marks the record at Planaltina, in Distrito Federal, and the white asterisk our records in south-east Goiás.

- R. J. (eds.) *The Cerrados of Brazil: ecology and natural history of a Neotropical savanna*. New York: Columbia University Press.
4. Pinto, O. M. O. (1936) Contribuição à ornithologia de Goyaz. *Rev. Mus. Paulista* 20: 1–171.
 5. Ridgely, R. S. & Tudor, G. (1994) *The birds of South America*, 2. Austin: University of Texas Press.
 6. Ridgely, R. S. & Tudor, G. (2009) *Field guide to the songbirds of South America: passerines*. Austin: University of Texas Press.
 7. Sick, H. (1997) *Ornitologia brasileira*. Rio de Janeiro: Ed. Nova Fronteira.
 8. Stotz, D. F., Fitzpatrick, J. W., Parker, T. A. & Moskovičs, D. K. (1996) *Neotropical birds: ecology and conservation*. Chicago: University of Chicago Press.
 9. Tubelis, D. P. (2008) First report on bird species collected by two Brazilian museums at Distrito Federal, central Cerrado. *Biotemas* 21: 157–161.
- Iubatã Paula de Faria**
PPG em Ecologia e Conservação, Universidade Federal do Mato Grosso do Sul, Campo Grande, MS, Brazil. Current address: QI 02, Bloco H, apto. 308, Guarã I, CEP 71010-080, Brasília, DF, Brazil. E-mail: iuba1@yahoo.com.br.
- Sandro Barata Berg and Tarcísio Lyra dos Santos Abreu**
Seriema Serviços e Estudos de Meio Ambiente, Brasília, DF, Brazil.
- Ana Paula Diniz Nakamura**
Graduação em Biologia, Universidade de Brasília, Brasília, DF, Brazil.
- Pedro Diniz**
Graduação em Biologia, Centro Universitário de Brasília, Brasília, DF, Brazil.

Received 18 February 2010; final revision accepted 25 September 2010 (published online 16 March 2011)